



Faculty of Women for Arts,
Science, and Education



Scientific Publishing Unit



Journal of Scientific Research in Science

Biological Sciences

Volume 38, Issue 1, 2021

ISSN 2356-8372 (Online) \ ISSN 2356-8364 (print)





Application of Gum Arabic as Edible Coating for Improving Postharvest Quality of Potato Tubers

Rowaa B. Ghannam^{a,*}, Shima M. Abdelsalam^b, Afaf A. Amine^a, Maha A. Hewedy^b

^aFood Safety Department, Microbiology Unit, National Nutritional Institute (NNI), Cairo, Egypt

^b Botany Department, Faculty of Women for Arts, Science and Education, Ain Shams University, Cairo, Egypt. Postal code 11757

Abstract

Hydrocolloid gums are extensively used in food industry. Recently hydrocolloid gums especially Gum Arabic (GA) has been widely used as edible coatings to extend shelf life of postharvest fruits and vegetables. The objective of this study was to evaluate the effect of GA edible film application mixed with glycerol and CaCl₂ as base matrix with all GA films i.e. GA/EDTA, GA/*L. paracasei* supernatant and GA/EDTA/*L. paracasei* supernatant in preservation of potato tubers stored at 8°C and 30±5°C for 35 days. Physicochemical analyses including pH, weight loss percentage and total soluble solids percentage (TSS), as well as microbial analysis (total counts of bacteria, mold and yeast and Enterobacteriaceae). The obtained result revealed that the total microbial count was found that GA/EDTA/*L. paracasei* supernatant gave a very close results to petroleum coating (wax) and the best results compared with noncoated (control) tubers, this was evident through the lower microbial load, better results than other treatments and noncoated tubers such as reducing weight loss, total soluble solids, better control on pH and expansion of the shelf life.

Keyword: Potatotubers, Postharvest quality, Edible coating, Gum Arabic, EDTA, *L. paracasei*, Microbial count.

1. Introduction

Potato (*Solanum tuberosum* L.) cultivation dates back to approximately 9000 years. Potato cultivation slowly spread throughout Europe in the following four centuries, and from there, to the rest of the world [22]. Potato by volume of production is the fourth important crop worldwide [46]. It is grown in about 140 countries; it is of high yielding and a low cost to the human diet and has a high nutritional value, as it contains about 40 IU vitamin A, 12 mg ascorbic acid, 77.8% water, protein, 2.1%, 0.3% fat, 22.6% carbohydrates, 1.1% crude fiber, 0.9% ash/100 g of edible portion. It is also used for production of high quality alcohol, starch, etc. and its starch (farina) is

***Corresponding author:** Rowaa B. Ghannam: Food Safety Department, Microbiology Unit, National Nutritional Institute (NNI), Cairo, Egypt.

E-mail: rowaaghannamm76@gmail.com

(Received 13 Mar 2021, revised 30 May 2021, accepted 8 Jun 2021)

<https://doi.org/10.21608/jsrs.2021.210680>

used in textile mills for sizing yarn in laundries. It is also used for the production of glucose and dextrin [23,46].

Edible coating is one of the promising techniques used in postharvest crops preservation [14]. Edible coating can be described as thin film of edible material that obstructs water transfer, gas exchange and soluble movement (i.e. semipermeable thin layer around the fruit which slow oxidation process, weight loss and respiration). It is safe, not synthesized chemically and can be an alternative to other traditional methods, such as chemicals and radiation that have undesirable effects on the human systems [16,38].

Gum Arabic is one of the hydrocolloid gums, it is water soluble secretion from Acacia tree branches especially *Acacia senegal* [15]. It is a highly valued gum because of its higher emulsifying, solubility, flavoring and thickening characteristics [4]. The prebiotic potential of Gum Arabic (GA) increases its health benefits, since it is rich in dietary fiber and indigestible in small intestine and stomach so it is fermented in large intestine by the surrounding bacteria [25,36,47]. Niamah *et al.* [42] reported that the viability of probiotic bacteria was increased when incorporated with GA.

Several newly methods for fresh fruits and vegetables microbial safety have been investigated, these methods were able to inhibit the colonization of pathogenic microorganisms in food products using Lactic Acid Bacteria (LAB), which have an antimicrobial capacity, since they have bacteriocins, lactic acid and ethanol and generally recognized as safe (GRAS) by the United States Food and Drug Administration (FDA), and thus makes the ideal for developing an environmentally friendly bioprotective agents in fresh fruits and vegetables [31,40]. Furthermore, LAB can be directly applied as food additives, or their purified metabolites or fermentation products can be used instead [19].

The spectrum of LAB can be successfully broadened when it is applied in combination with chelating agent as Ethylene diamine tetra-acetic acid (EDTA)[39]. EDTA well-known chelating agent has been widely employed to control oxidation and other metal ion-catalyzed deteriorative reactions in food products. EDTA disodium or calcium disodium salts have been approved as food additives by FDA [19]. It also preserves color, flavor and odor and it is worth mentioning that it doesn't accumulate in biological tissues and is not carcinogenic in 500mg/kg/day volume [7]. According to literature data, EDTA is known antimicrobial compounds [18], what was confirmed by

Borowicz *et al.* [10], where EDTA has antimicrobial activity against some gram positive and gram negative bacteria.

Calcium is a key plant nutrient that has a significant role in cell functions, at various concentrations, as delaying the ripening, softening and senescence [9], also cross linking with calcium chloride generates stronger film gels [29].

Meeting the criteria of high quality and reduce the use of chemicals to obtain food safety products are few of the factors contributing to this requirement. This has lead researchers and companies to explore different ways to improve their productivity in terms of maintaining freshness, quality and food safety, such as using sustainable materials in food packaging. Thus this study aims to evaluate the effect of glycerol (as plasticizer), CaCl₂ and GA with both EDTA and *L. paracasei* cell free supernatant in the component of potato edible film/coating and their effect on the pathogenic microbial composition and load in compared with wax as a petroleum product, to prolong the shelf life of potato tubers for local markets and exportation.

2. Materials and Methods

2.1 Sample collection:

Three groups of potato tuber samples were collected from local market, Cairo, Egypt. They were transported to Microbiology Unit, Food Safety department, National Nutritional Institute (NNI), Cairo, Egypt. The experiments were conducted for approximately 35 days under cold (8°C) and ambient temperature (30±5°C). The first group was let to spoil for isolating the spoiled microorganisms that affect potato crops. The second group was selected as fresh, mature stage and homogenous appearance based on color, size and free of mechanical damage. The third group was prepared for coating by washing with tap water (5 mg/L free chlorine) [12] and dried at ambient air conditions before coating for proper adherence to the surface.

2.2 Bacteria strain:

Lactobacillus Paracasei isolate was obtained from National Center for Radiation Research and Technology (NCRRT), Egyptian Atomic Energy Authority (EAEA), Cairo, Egypt.

Stock culture preparation and growth condition of *L. paracasei*:

Lactic acid bacterial isolate; *L. paracasei* was stored at -80°C MRS broth supplemented with 30% v/v sterile glycerol, it was reactivated and recultured under anaerobic conditions in CO_2 incubator (BPI, Italy) at 37°C for 48h. Subsequently, appropriate serial dilutions were prepared and precultured at plate count agar medium (PCA, Oxoid), to obtain 10^6 CFU/ml of *L. paracasei* [27,35]. The colonies were harvested with a sterilized loop and inoculated into screw-capped tubes of 9ml MRS broth, then centrifuged at 4000rpm for 30 minutes at 4°C [45]. The cell free supernatant was separated and the pellet was discarded.

2.3 Coating treatment

2.3.1 Gum Arabic (GA) edible film

Ten g GA obtained from Sudan (from local market) was dissolved in 100ml distilled water and heated at 60°C for 30min. [2,17] with gentle stirring, and then it was filtered. For all edible films preparation after cooling to 50°C , glycerol of (1.0% v/v) (99% purity, Sigma) as a plasticizer to improve the flexibility of coating solutions and CaCl_2 of (1.0% w/v) were added to GA solution [32].

The GA edible film was divided into four groups, one group was for the film only, the second was added to 2.5% Calcium disodium Ethylene Diamine Tetraacetate salt (EDTA), (Sigma, Aldrich- Germany) [10], the third group was GA film with *L. paracasei* supernatant, (50% w/v) [27], and the fourth one was for the film in combination with both EDTA and *L. paracasei* supernatant.

2.3.2 Wax film

Commercial Paraffin wax was melted and prepared for covering film

2.3.3 Application of coating solutions

The films were applied to the tuber samples by a casting method while the wax film was applied with a brush. All coated and noncoated tuber samples (control) were divided into two groups, one stored at 8°C and the other at $30\pm 5^{\circ}\text{C}$. During the storage period microbial analysis were performed every 3, 5, 7, 10, 15, 20, 25, 30 and 35 days.

2.4 Microbial analysis

Both coated and noncoated crop samples were peeled off and 2cm^2 of the inner tissue was transferred for microbial examination to screw-capped tube with 9 ml buffered peptone water (0.1%) under sterile conditions and vortexed for 2min.

Appropriatedilutions were prepared for the microbial analysis.The microbial analysis consisted of a plate count method for bacteria at 37°C for 24h in a plate count agar (PCA), mold and yeast at 22-25°C for 5 days in Sabouraud (SB) agar and enterobacteriaceae at 37°C for 24h in double layer Violet Red Bile Glucose Agar. Visible colonies were counted and CFU/g calculated. All analysis were performed in triplicates.

2.4 Postharvest quality

2.4.1 pH determination

Five grams of groundedpotato tuber samples for both coated and noncoated samples in 50 ml distilled water were used to measure samples pH at 25°C with subsequent reading in a digital pH- meter (240A Orion, USA).

2.4.2 Weight loss percentage

Weight loss (expressed as percentage) for both coated and noncoated tubers were calculated from zero time and days 3, 5, 7, 10, 15, 20, 25, 30, 35 during the storage period, using the following equation:

$$\text{Weight loss} = \frac{\text{fruit initial weight} - \text{fruit weight on sampling day}}{\text{fruit initial weight}} \times 100 \text{ [32].}$$

2.4.3 Total soluble solids TSS percentage

During the storage, the TSS percentage was determined from grounded (as mentioned in pH) tuber samples, using Car Zeiss hand refractometer, Germany, as described in AOAC [5].

2.5 Scanning Electron Microscopy (SEM)

The outer peel of both coated and noncoated potato tuber samples of the promising film (GA/EDTA/*L. paracasei* supernatant) were fixed in 2-5% glutaraldehyde for 24h. at 4°C and was post fixed in 1.0% osmium tetroxide for 1h. at room temperature [24]. Then the specimens were dehydrated with ascending concentrations of acetone, critical point dried, and finally sputter was coated with gold. The photographing and examination had been done through Jeol Scanning Electron Microscope (JSM-5200LV-LGS, JEOL Japan) and accelerating voltage of 30kv and magnified 50 and 100X.The SEM was performed at Applied Nematode Incest Centre, Faculty of Agriculture, Cairo University.

2.6 Statistical analysis

The experiments were carried out in completely randomized design with two replicates. Analysis of variance (ANOVA) was conducted according to Casella [11], using Statistix 10 software program for discriminating the treatment means, the Duncan's multiple range test at $P < 0.05$ was used

3. Results

Results in Tables (1&2) representing the effect of GA edible films and wax on total Enterobacteriaceae count of both coated and noncoated potato tubers in both cold and ambient temperatures, data showed better control of GA/EDTA/*L. paracasei* supernatant film as well as wax on tubers compared with noncoated (control) tubers. During 35 days of storage the total Enterobacteriaceae count for all treated tubers was ≤ 10 CFU- g^{-1} while control samples recorded counts after twenty days at cold storage and represented as 10 CFU- g^{-1} , day 25 represented as 70 CFU- g^{-1} and day 30 (deterioration day) recorded 300 CFU- g^{-1} . Approximately same results were detected at ambient storage temperature, from day 15 and represented as 80 CFU- g^{-1} and recorded 700 CFU- g^{-1} at deterioration day (day 20).

Table (1): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on the total Enterobacteriaceae count during storage at 8°C

Storage days Treatments	0	3	5	7	10	15	20	25	30	35
Gum Arabic	≤10	≤1 0	≤1 0	≤10	≤10	≤10	≤10	≤10	≤10	≤10
Gum Arabic/ EDTA	≤10	≤1 0	≤1 0	≤10	≤10	≤10	≤10	≤10	≤10	≤10
Gum Arabic/ <i>L. paracasei</i> supernatant	≤10	≤1 0	≤1 0	≤10	≤10	≤10	≤10	≤10	≤10	≤10
Gum Arabic/ EDTA/ <i>L. paracasei</i> supernatant	≤10	≤1 0	≤1 0	≤10	≤10	≤10	≤10	≤10	≤10	≤10
Wax	≤10	≤1 0	≤1 0	≤10	≤10	≤10	≤10	≤10	≤10	≤10
nonncoated Control	≤10	≤1 0	≤1 0	≤10	≤10	≤10	≤10	70	300	cd

cd =complete deterioration

Table (2): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on the total Enterobacteriaceae count during storage at 30±5°C

Storage days	3	5	7	10	15	20	25	30	35
Treatments									
Gum Arabic	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10
Gum Arabic/ EDTA	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10	cd
Gum Arabic/ <i>L. paracasei</i> supernatant	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10	0
Gum Arabic/ EDTA/ <i>L. paracasei</i> supernatant	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10	0
Wax	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10	≤10
nonncoated (Control)	≤10	≤10	≤10	≤10	80	700	cd	cd	cd

cd =complete deterioration

On examining the ability of GA/EDTA/*L. paracasei* supernatant edible film to preserve potato tubers from rotting, rooting, shrinking and sprouting compared with wax film and noncoated tubers at cold storage results in Figure (1) were obtained.




Fig. (1): photo of (a) noncoated potato tuber, (b) coated tubers with GA/EDTA/*L.paracasei* supernatant film and (c) coated tubers with wax film after 35 days at 8°C.




Fig. (2): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on bacterial, yeast and fungal count during the storage at 8°C (a: total bacterial count, b: yeast, c: fungi) and at 30±5°C (d: total bacterial count, e: yeast, f: fungi).

Results in Figure (2a,b and c) showed that the cold storage (8°C) of potato tubers that coated with GA, GA/*L.paracasei* supernatant, GA/EDTA/ *L. paracasei* supernatant extend the shelf life of tubers as well as wax more than 35 tested days and tubers coated with GA/EDTA edible film were spoiled at day 35 and all films prevented spoilage compared with noncoated tubers (control), that decayed within 25 days, while Figure (2d,e and f): at 30±5°C storage the samples coated with GA, GA/*L. paracasei* supernatant, GA/EDTA/ *L. paracasei* supernatant and wax continued up to 30 days without decay, while the tubers that coated with GA/EDTA were deteriorated at day 30 and bacterial and yeast count were recorded, although the coated tubers were better than noncoated (control) samples since the bacterial count and yeast and fungal counts were clear in addition to the complete decaying at day 20. As for pH, the increment of storage period led to a slight increase in pH value of stored potato tubers (Tables 3&4), especially at the end of storage period. The noncoated tubers scored lower pH value than other treatments and wax coating. There were varieties between different treatments especially tubers coated with GA/EDTA/*L. paracasei* supernatant edible film that had the highest value of pH as well as wax coating during both the low and ambient storage temperature compared with other edible films and noncoated tubers which scored lower pH value.

Table (3): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on pH during the storage at 8°C

Treatments	Storage time/ days										
	0	3	5	7	10	15	20	25	30	35	mean
Gum Arabic	5.84	5.92	6.18	6.33	6.29	6.15	6.2	6.26	6.3	6.33	6.20 ^{bc}
Gum Arabic+EDTA	5.79	5.83	5.99	6.19	6.23	6.27	6.3	6.35	6.37	6.41	6.19 ^{bc}
Gum Arabic+<i>L. paracasei</i> supernatant	5.82	5.86	5.93	6.2	6.09	6.46	6.62	6.65	6.66	6.69	6.31 ^b
Gum Arabic+EDTA+<i>L. paracasei</i> supernatant	5.81	5.95	6.08	6.26	6.39	6.57	6.7	6.73	6.75	6.78	6.43 ^b
Wax	5.84	6.05	6.27	6.63	6.85	6.99	7.1	7.12	7.16	7.17	6.77 ^a
control	5.8	5.83	5.87	5.92	5.94	5.97	6	6.02	6.05	cd	5.97 ^c
mean	5.82 ^d	5.91 ^{cd}	6.05 ^{bcd}	6.255 ^{abc}	6.30 ^{abc}	6.40 ^{ab}	6.48 ^{ab}	6.62 ^a	6.65 ^a	6.61 ^a	

Means followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

cd= complete deterioration.

Table (4): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on pH during the storage at 30±5°C

Treatments	Storage time/ days									
	0	3	5	7	10	15	20	25	30	mean
Gum Arabic	5.78	5.8	5.84	5.94	6.3	6.55	6.89	6.9	6.91	6.32 ^{ab}
Gum Arabic+EDTA	5.9	6.3	6.43	6.81	5.93	6.16	6.22	6.43	6.57	6.31 ^b
Gum Arabic+<i>L. paracasei</i> supernatant	5.89	6.15	6.32	6.55	6.16	6.09	6.21	6.79	6.84	6.33 ^{ab}
Gum Arabic+EDTA+ <i>L. paracasei</i> supernatant	5.80	5.83	5.92	6.17	6.30	6.43	6.62	7.03	7.05	6.35 ^{ab}
Wax	5.86	6.20	6.37	6.73	6.86	6.92	6.99	7.01	7.02	6.66 ^a
Control	5.86	5.89	5.92	5.98	6.01	6.12	6.15	cd	cd	6.12 ^b
mean	5.85 ^c	6.03 ^{cd}	6.13 ^{bcd}	6.36 ^{abc}	6.26 ^{bcd}	6.38 ^{abc}	6.51 ^{ab}	6.77 ^a	6.83 ^a	

Means followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

cd= complete deterioration.

Tables (5) and (6), show a gradual increasing in weight loss percentage for all samples during both low and ambient storage temperature, while the tubers coated with wax and GA/EDTA/*L. paracasei* supernatant edible film scored a significant lowest weight loss percentage compared with tubers coated with other treatments or noncoated ones.

Table (5): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on percentage of weight loss during the storage at 8°C

Treatments	Storage time/ days									
	3	5	7	10	15	20	25	30	35	mean
Gum Arabic	0.19	0.25	0.42	0.67	0.82	1.01	1.13	1.29	1.34	0.72 ^{bc}
Gum Arabic+EDTA	0.21	0.33	0.48	0.72	1.02	1.15	1.26	1.3	1.31	0.81 ^b
Gum Arabic+<i>L. paracasei</i> supernatant	0.12	0.18	0.34	0.61	0.88	0.96	1.2	1.25	1.28	0.69 ^{cd}
Gum Arabic+EDTA+ <i>L. paracasei</i> supernatant	0.1	0.15	0.3	0.57	0.79	0.99	1.06	1.12	1.21	0.64 ^d
Wax	0.06	0.11	0.27	0.4	0.55	0.67	0.72	0.77	0.81	0.44 ^e
Control	0.39	0.65	0.98	1.13	1.37	1.63	1.88	1.89	cd	1.30 ^a
mean	0.17 ^f	0.27 ^f	0.47 ^e	0.68 ^d	0.91 ^c	1.07 ^b	1.07 ^a	1.15 ^a	1.29 ^a	

Means followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

cd= complete deterioration.

Table (6): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on percentage of weight loss during the storage at 30±5°C.

Treatments	Storage time/ days								
	3	5	7	10	15	20	25	30	mean
Gum Arabic	0.21	0.31	0.64	0.89	1.36	1.49	1.7	1.75	1.04 ^{bc}
Gum Arabic+EDTA	0.25	0.36	0.66	0.92	1.38	1.51	1.71	1.77	1.07 ^b
Gum Arabic+<i>L. paracasei</i> supernatant	0.19	0.29	0.62	0.88	1.33	1.47	1.69	1.74	1.03 ^{bc}
Gum Arabic+EDTA+<i>L. paracasei</i> supernatant	0.17	0.25	0.59	0.93	1.28	1.43	1.65	1.69	1.00 ^c
Wax	0.15	0.22	0.5	0.82	1.11	1.25	1.32	1.38	0.84 ^d
Control	0.46	0.775	1.1	1.47	1.83	1.95	cd	cd	1.48 ^a
mean	0.24 ^g	0.37 ^f	0.69 ^e	0.99 ^d	1.39 ^c	1.51 ^b	1.69 ^a	1.74 ^a	

Means followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

cd= complete deterioration.

Total soluble solids percentage (TSS%) of potato tubers was presented in Tables (7) and (8), it was increase proportionally with the increase of storage period for both storage temperatures, where treated tubers significantly retained lower TSS than the control. These results revealed that GA/EDTA/*L. paracasei* edible film supernatant and wax coatings efficiently reduced the rapid changes in TSS.

Table (7): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on percentage of Total Soluble Solids during the storage at 8°C

Treatments	Storage time/ days										
	0	3	5	7	10	15	20	25	30	35	mean
Gum Arabic	5.75	5.79	5.87	6.17	6.26	6.33	6.29	6.17	6.19	6.22	6.10 ^c
Gum Arabic+EDTA	6.25	6.26	6.38	6.46	6.49	6.34	6.48	6.5	6.53	6.57	6.43 ^b
Gum Arabic+L. paracasei supernatant	5.81	5.83	5.95	6.09	6.17	5.96	6.07	6.2	6.23	6.38	6.07 ^c
Gum Arabic+EDTA+ L. paracasei supernatant	5.72	5.75	5.78	5.82	5.83	5.89	5.7	5.69	5.66	5.63	5.75 ^d
Wax	5.63	5.64	5.65	5.73	5.74	5.81	5.75	5.71	5.73	5.78	5.72 ^d
Control	6.5	6.54	6.66	6.89	7.12	7.2	7.03	6.99	6.95	cd	6.88 ^a
mean	5.94 ^a	5.97 ^a	6.05 ^a	6.19 ^a	6.27 ^a	6.255 ^a	6.22 ^a	6.21 ^a	6.22 ^a	6.26 ^a	

Means followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

cd= complete deterioration.

Table (8): The effect of Gum Arabic edible films, wax film and noncoated (control) potato tubers on percentage of Total Soluble Solids during the storage at 30±5°C

Treatments	Storage time/ days									
	0	3	5	7	10	15	20	25	30	mean
Gum Arabic	5.53	5.56	6.01	6.51	5.95	6.13	6.32	6.4	6.44	6.09 ^{ab}
Gum Arabic+EDTA	5.62	5.83	6.25	6.25	6.63	6.07	6.18	6.22	6.3	6.15 ^{ab}
Gum Arabic+<i>L. paracasei</i> supernatant	5.7	5.77	5.84	5.91	6.17	6.22	6.18	6.2	6.07	6.01 ^{ab}
Gum Arabic+EDTA+<i>L. paracasei</i> supernatant	5.67	5.69	5.79	5.85	5.82	6.09	6.22	6.34	6.35	5.98 ^{ab}
Wax	5.43	5.45	5.6	5.83	5.93	6.18	6.25	6.33	6.36	5.93 ^b
Control	5.67	5.89	6.03	6.15	6.25	6.63	6.85	cd	cd	6.29 ^a
mean	5.60 ^c	5.70 ^{bc}	5.92 ^{abc}	6.08 ^{ab}	6.13 ^{ab}	6.22 ^a	6.33 ^a	6.34 ^a	6.30 ^a	

Means followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test

cd= complete deterioration




Figure (3): SEM photo of (a) noncoated potato tubers, (b) coated potato tuber with GA/EDTA/*L. paracasei* supernatant

Figure (3 a&b) represents pictures of potato tubers' peel which were captured for the best edible film GA/EDTA/*L. paracasei* supernatant by Scanning Electron Microscope (SEM) to study the physical featured of coated potato tubers compared with thenoncoated tubers. Figure (3a) represents noncoated tuber peel, which has district separation, while Figure (3b) showed homogeneity, minimum pores and cracks and better structural continuity of edible coating.

4. Discussion

Food safety and quality is the most important standard for every consumer and also it is the priority for food industry. Edible films/coatings provide edibility, biodegradability, compatibility and eco-friendly food food packaging option instead of petroleum option like wax, especially for ready to eat food like vegetables and fruits.

Contamination of food with spoiled microorganisms reduces its quality; however the existing pathogenic microorganisms present a serious health problem. Temperature, humidity, pH, water as well as carbohydrate content of potato tubers greatly affect the behavior of pathogens. Lazarovits *et al.* [28] reported that about 10^7 CFU/g of the soil lived in potato rhizosphere and geocaulosphere; which is the volume of the soil surrounding the tubers.

Temperature and humidity in general provide a favorable environment for the development of microorganisms. Therefore we investigated the ability of GA, GA/EDTA, GA/ *L. paracasei* supernatant and GA/EDTA/*L. paracasei* supernatant

edible film supernatant/coatings to prevent microbial development in potato tubers both in cold (8°C) and in an ambient temperature (30±5°C). In fact GA/EDTA/*L. paracasei* supernatant coating showed a remarkably inhibitory effect on microbial growth during storage period where cold storage (8°C) helps to extend shelf life of tubers up to 35 days than ambient temperature (30±5°C) that preserve tubers up to 30 days, as shown in Figure (2a,b,c,d,e and f). This synergistic effect against spoiled strains is enhanced by hycolloidal edible coatings such as Gum Arabic that has a protective effect since it acts as a barrier to moisture and gas, also it has antioxidant action [41], and antimicrobial activity against some strains of G-ve, G+ve bacteria, fungi and yeasts [30].

The prepared film (Gum Arabic+EDTA+ *L. paracasei* supernatant) was found to prevent both rooting and sprouting growth at 8°C as shown in Figure (1), since storage of potato tubers at 8–12 °C is common practice, where the accumulation of sugars is minimum and potatoes remain suitable for processing. However, this temperature is suitable for sprout growth. Therefore, use of a sprout suppressant becomes essential. Isopropyl N-(3-chlorophenyl) carbamate (CIPC) is the most commonly used sprout suppressant used in potatoes. This is the most common way of long-term (up to 6 to 9 months) storage of potatoes in tropical, sub-tropical and sub-temperate countries of the world. CIPC belongs to group of pesticides known as carbamates, 3-chloroaniline (3-CA) this is considered more polluting and highly toxic to human and environment. CIPC is more harmful to birds, fishes and other aquatic animals, hence its washing contaminate water bodies, environment and worms. There are growing concerns regarding degradation products/metabolites of CIPC as they are more toxic and cytolytic in nature; highly toxic, carcinogenic, cause reduction in ATP synthesis so bring about modifications in cell permeability besides being pollutants [43]. Hence, Application of natural products that applied in the present study may help in solving such problems.

Saha *et al.* [50] investigated the effect of chitosan and whey protein in the quality of potato tubers and extend its shelf life to 60 days at 20°C. Also Marquez *et al.* [37] preserved potato tubers up to 10 days by whey protein/ pectin/ transglutaminase coating that show a significant inhibitory effect on mesophilic colonies growth in all tuber samples. Calcium chloride helps to generate stronger film gel as it affects the kinetics of gelation, so the steady-state gel strength was reached fastest by CaCl₂ [29].

CaCl₂ has significant role in plant cell function at various concentrations as delayed the ripening, softening and senescence [18]. It has been also reported to aid in reduction of browning by inhibiting poly phenol oxidase (PPO) which has been demonstrated for potato [8]. The presence of LAB in the matrix that produce antimicrobial substances such as bacteriocins or organic acid, it also decreases pathogen adhesion, promote recovery or enhance stability of commensal microbiota when perturbed and delivered functional enzymes proteins as reported by Gayathri and Ramesha [21], this is supported with Maarof *et al.* [35], who concluded the antimicrobial of probiotic strains isolated from milk products against some common food-borne pathogens as they show bactericidal effect on *E. coli* and *Salmonella spp.* while have bacteriostatic effect on *Staphylococcus aureus.* and against *R. solanacearum.*

Washing with tap water that contains 5mg/L free chlorine has significant effect on spoilage microorganisms. Similar results was obtained by the work of Lu *et al.* [34], who found that the final bacterial number predicted on lettuce treated with chlorinated water was lower than that on non-treated lettuce. This also agrees with that obtained by Rico *et al.* [49], who found that washing with chlorinated water decontaminate vegetables.

EDTA supports and potentiates the sensitivity of Gram negative bacteria to bacteriocin particularly nisin; a propiotics' product [34]. EDTA inhibits bacterial adhesion as it has antimicrobial properties against *P. flourescens*, *E. coli*, *Bacillus sp* and *Proteus sp.*[10]. Trejo-Gonzalez *et al.* [51] also reported its inhibitory effect.

On the other hand the presence of EDTA may causea reduction in GA antimicrobial activity; several mechanisms could be responsible for the antagonistic effect between GA and EDTA. EDTA might prevent GA from being taken up by the cell or vice versa. Alternatively, the interaction between EDTA and GA may prevent GA from reaching its target of its activity or it mayinterfere with its composition or structure its specific activityto be reduced this need further investigation.

The active food packaging include desirable interactions with food, in a way that is related to extending food stability, by interacting with product or headspace inside to prevent, decrease or delay microbial growth onthe food surface [15], in contrast to wax which act as barrier possesses no interaction with theproduct. This gasses exchange barrier blocking the supply of O₂ through the tuber peel needed for

respiration or CO₂ releasing, this block causes off-flavor development due to anaerobic fermentation and this is in agreement with the results of previous studies [52].

For pH value in this study, it was noticed that the progress in storage period slightly increases the pH value, the same finding was found by Saha *et al.* [50] on stored potato tubers by chitosan. GA/EDTA/*L. paracasei* supernatant edible film showed better control on pH as well as wax coating during entire storage periods as compared either to other treatments or to the noncoated potato tuber this might be referred to more moisture content, increasing respiration rate and the conversion of organic acids to sugars, which is consumed during storage in some metabolic processes [27].

Weight loss is a quality index of the post-harvest life of tubers. Weight loss of treated potato tubers was significantly lower than untreated tubers during the entire storage periods as represented in Tables (5 and 6), the similar results was reported by Khaliq *et al.* [26], in which weight loss was reduced by GA treatment for mango fruits. This reduction in fresh weight of tubers was attributed to the effect of edible coating that act as a fence against gas exchange and water loss through respiration and transpiration [1,37]. In contrast to noncoated tubers, the increase in weight loss is due to the increase in transpiration process and cellular breakdown of crop samples [13].

Total soluble solids (TSS) percentage is a maturity and quality measurements of tubers. In this study, TSS gradually increased in all samples regardless of treatments as the storage time increase. Similar trend in the results was revealed in mango [31,53] and also previous report by Saha *et al.* [50] on potato tubers. The slight increasing in TSS is due to hydrolysis of potato starch by the action of enzymes into sugars. GA act as barrier to gas exchange, results in delaying the respiration rate and hence hydrolytic enzymes action. This is reflected on carbohydrate consumption that decreasing TSS content and this agrees with Khaliq *et al.* and El-abbasy *et al.* [13,26].

Photos (Figures 3: a&b) of potato tuber peel were captured by scanning electron microscope (SEM) to study the physical structure of coated tubers with the best film, compared to noncoated (control). Figure (3a) represents the noncoated tuber peel which has district separation, while Figure (3b) shows better structure content, homogenization and minimum pores and cracks compared with control. This structure reflects the gelatinized GA matrix dissolved in *L. paracasei* supernatant edible film, that was homogenous but not heavy and the presence of glycerol (plasticizer) provides flexibility, well thickness for the film in addition to a good moisture content, this was

agreed with Adjoumar *et al.* [3] in cassava film that was homogenous, crack free, transparent due to glycerol content.

5. Conclusion

From this study we can conclude that the combination of Gum Arabic, EDTA and *L. paracasei* supernatant edible film can be used as an alternative choice for petroleum products (wax) to keep potato tubers as a safe, environmentally friendly, and in reducing microbial load in comparison with petroleum product as wax.

6. Acknowledgement

We are thankful for National Nutritional Institute and Faculty of women for Arts, Science and Education Ain Shams University for offering lab equipments and their technical support.

7. Conflict of Interest

The authors have declared no conflict of interest.

References

- [1] A.F. Abd El-khalek,, Effectiveness of gum arabic, potassium salts and their incorporation in the control of postharvest diseases and maintaining quality of ‘Washington’ navel oranges during long term cold storage, *Egyptian Journal of Horticulture*45 (2): (2018)185-203
- [2] A. Ali, M. Maqbool, S.Ramachandran, and P. G. Alderson,Gum arabic as a novel edible coating for enhancing shelf-life and improving postharvest quality of tomato (*Solanum lycopersicum* L.) fruit, *Postharvest biology and technology* 58(1): (2010) 42-47
- [3] Y. D. Adjouman, C.Nindjin, K. N. Kouassi, F. A.Tetchi, G. A. N'Guessan, and M. Sindic, Effect of edible coating based on improved cassava starch on post-harvest quality of fresh tomatoes (*Solanum lycopersicum* L.), *International Journal of Nutritional Science and Food Technology* 4(1) :(2018) 1-10.
- [4] C. Amine, V. Mejia Tamayo, T. Doco, C. Sanchez, M. Nigen, P. Williams, and D. Renard, Acacia gum: History of the future, *Food Hydrocolloids* 78: (2017) 140–160
- [5] AOAC 1990 Official methods of analysis (15th ed.) Arlington, VA: Association of Official Analytical Chemists. Atanasova, J., Moncheva, P., & Ivanova, I. (2014). Proteolytic and antimicrobial activity of lactic acid bacteria grown in goat milk. *Biotechnology and Biotechnological Equipment* 28:1073–1078
- [6] S. Atrash, A. Ramezani, M. Rahemi, R. M. Ghalamfarsa, and E. Yahia, Antifungal effects of savory essential oil, gum arabic, and hot water in Mexican lime fruits, *Hort Science* 53: (2018)524–530
- [7] N. Bagheripoor, S. Khoshgozaran-Abras, S. Sohrabvandi, N. Khorshidian, A. M. Mortazavian, N. MollaKhalili, and S. Jazaeri, Application of Active Edible

- Coatings to Improve the Shelf-life of Cheese, Food Science and Technology Research, 24(6): (2018) 949-962
- [8] E. A. Baldwin Coatings and other supplemental treatments to maintain vegetable quality,(2003), Marcel Dekker, New York. p. 413
- [9] A.V. Barker, and D.J. Pilbeam, Handbook of Plant Nutrition, (2015), Florida 773
- [10] J. Borowicz, K. Kasperski, D. Gwiazdowska, M. Tichoniuk, P. Wojciechowska, and K. Juś, " Antibacterial activity of chemical compound used for packaging" (2016), <http://ue.poznan.pl>
- [11]V. Dhundale, V. Hemke, D. Desai, and P. Dhundale, Evaluation and exploration of lactic acid bacteria for preservation and extending the shelf life of fruit, International Journal of Fruit Science (2018)1–16.
- [12] Egyptian Minister of Health and Population decision No. 204 of 2007 regarding the maximum limits of standards and specifications that must be met in safe drinking water and domestic use
- [13] U. El-Abbasy, A. El-khalek, H. Maswada, and A. Abou-Ismaïl, Responce of physival and physio-chemical attributes in Guava fruits Gum Arabic and calcium chloride treatments under cold storage, Journal of Productivity and Development, 24(4): (2019) 705-725.
- [14] P. J. P. Espitia, W.X. Du, R. De Jesús Avena-Bustillos, N. F. F. D. Soares, and T. H. McHugh, Edible films from pectin: Physical-mechanical and antimicrobial properties - A review, Food Hydrocolloids 35: (2014) 287–296
- [15] P. J. Espitia, R. A. Batista, H. M. Azeredo, and C. G. Otoni, Probiotics and their potential applications in active edible films and coatings. Food Research International, 90: (2016). 42-52.
- [16] M. M. Falcao-Rodrigues, M. Moldao-Martins, and L. M. Beirão-dacosta, DSC as a tool to assess physiological evolution of apples preserved by edibles coatings, Food Chemistry 102: (2007) 475–480
- [17] T. A. Fashanu, A. T. Oladiji, and O. A. Peters, Efficacy of Gum Arabic as an esculent film on shelf life extension of tomato (*Solanum lycopersicum* L) fruit, Ruhuna Journal of Science 10(1): (2019) 32-50
- [18] S. Finnegan, and S. L. Percival, EDTA:an Antimicrobial and Antibiofilm Agent for Use in Wound Care, Advances in wound care 4(7): (2015) 415-421
- [19] Y. O. García, C. E. S. Maldonado, and M. Quirasco, Detección de proteínas con actividad antibacteriana producidas por bacterias ácido lácticas, Biotecnología 19:(2015) 25–43
- [20] W. E. Gawad, O. M. Helmy, W. M.,Tawakkol, and A. M. Hashem, Effect of EDTA on biofilm formation and antibiotic susceptibility of multidrug resistant uropathogenic Escherichia coli clinical isolates in Egypt, African Journal of Microbiology Research 11(38): (2017) 1445-1458
- [21] D. Gayathri, and A. Ramesha, Gluten-hydrolyzing probiotics: An emerging therapy for patients with celiac disease, World Academy of Sciences Journal 2(5): (2020) 1-1
- [22] C. Gebhardt, The historical role of species from the *Solanaceae* plant family in genetic research, Theoretical and Applied Genetics 129(12): (2016) 2281-2294.

- [23] D. Hada, and K. Sharma, A review: plant extracts a new source of antimicrobial potential, *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 5(3): (2014) 597-628.
- [24] M. M. Harley, and I. K. Ferguson, The role of the SEM in pollen morphology and plant systematics. *Scanning electron microscopy in taxonomy and functional morphology*, Oxford University Press, Oxford, (1990) 45-68.
- [25] O.H.M. Idris, What is gum arabic? an overview, *International Journal of Sudan Research* 7(1): (2017) 1–14.
- [26] G. Khaliq, M.T.M. Mohamed, P. Ding, H.M. Ghazali, and A. Ali, Storage behaviour and quality responses of mango (*Mangifera indica* L.) fruit treated with chitosan and gum arabic coatings during cold storage conditions, *International Food Research Journal*, 23: (2016) p.S141.
- [27] O. A. Khalil, A. M. Mounir, and R. A. Hassanien, Effect of gamma irradiated *Lactobacillus* bacteria as an edible coating on enhancing the storage of tomato under cold storage conditions, *Journal of Radiation Research and Applied Sciences* 13(1): (2020) 317-329
- [28] G. Lazarovits, J. Hill, G. Patterson, K. L. Conn, and N. S. Crump, Edaphic soil levels of mineral nutrients, pH, organic matter, and cationic exchange capacity in the geocaulosphere associated with potato common scab, *Phytopathology* 97(9): (2007) 1071-1082.
- [29] P. Lee, and M.A. Rogers, Effect of calcium source and exposure-time on basic caviar spherification using sodium alginate, *Int. J. Gastron Food Sci.* 1: (2012) 96-100.
- [30] P. Li, X.Zheng, Y.Liu, and Y.Zhu, Pre-storage application of oxalic acid alleviates chilling injury in mango fruit by modulating proline metabolism and energy status under chilling stress, *Food Chemistry* 142: (2014) 72–78.
- [31] D. Li, K. Ni, , H. Pang, Y. Wang, Y. Cai, and Q. Jin, Identification and antimicrobial activity detection of lactic acid bacteria isolated from corn stover silage, *Asian- Australasian Journal of Animal Science* 28:(2015) 620–631.
- [32] S. Licodiedoff, L.A. Koslowski, L. Scartazzini, A.R. Monteiro, J.L. Ninow, C.D. Borges, Conservation of physalis by edible coating of gelatin and calcium chloride, *International Food Research Journal* 23: (2016) 4
- [33] J. P. Lima da Silva, E. F. de Souza, R. C. Della Modesta, I. A. Gomes, O.Freitas-Silva, and B. D. Gombossy de Melo Franco, Antibacterial activity of nisin, oregano essential oil, EDTA, and their combination against *Salmonella* Enteritidis for application in mayonnaise, *Vigilancia Sanitaria Em debate-Sociedade Ciencia & Tecnologia* 4(1): (2016) 83-91
- [34] Z. Lu, L. Zhang, F. Lu, X. Bie, and Z. Yu, Model of microbial growth on fresh cut lettuce treated with chlorinated water during storage under different temperature, *Journal of Food Process Engineering* 29: (2006) 106–118.
- [35] H.A. Maarof, M.I.M. Abdallah, M.S. Bazalou, and R.G.Abo-Samra, (2013), "Effect of Probiotics Bacteria Isolated from Yoghurts Produced in Damietta City on Some Pathogenic Bacteria", *Proc. of the 6th Animal Wealth Research Conf. in the Middle East & North Africa* 1-14.

- [36] A.A. Mariod, (Ed.), (2018), "Functional Properties of Gum Arabic. In Gum Arabic: Structure, Properties", Application and Economics, p. 283–295. United Kingdom: Elsevier Inc. <https://doi.org/10.1016/B978-0-12-812002-6.00024-5>
- [37] G. R. Marquez, P.Di Pierro, L.Mariniello, M. Esposito, C. V. Giosafatto, and R.Porta, Fresh-cut fruit and vegetable coatings by transglutaminase-crosslinked whey protein/pectin edible films, LWT 75:(2017) 124-130
- [38] D. Martínez-Romero, N. Albuquerque, J. Valverde, F. Guillén, S. Castillo, D. Valero, and M. Serrano, Postharvest sweet cherry quality and safety maintenance by *Aloe vera* treatment: A new edible coating, Postharvest Biology and Technology 39: (2006).93–100.
- [39] M. A. Montenegro, M. L. Boiero, L.Valle, and C. D. Borsarelli, Gum Arabic: more than an edible emulsifier, Products and applications of biopolymers 51: (2012) 953-978
- [40] J. R. L. Morales, N. G. Méndez, B. E. R. Chavira, S. B. P. Vega, and G. V. N. Moorillón, Biocontrol processes in fruits and fresh produce, the use of lactic acid bacteria as a sustainable option, Frontiers in Sustainable Food Systems (2018) 1–13.
- [41] M. E. Mirghani, A. A. Elnour, N.Kabbashi, M. Z.Alam, , K. H. Musa, and A.Abdullah, Determination of antioxidant activity of gum arabic: An exudation from two different locations, Sci. Asia 44: (2018) 177-186.
- [42] A.K. Niamah, S.T.G. AL-Sahlany, and A.J. Al-Manhel, Gum Arabic Uses as Prebiotic in Yogurt Production and Study Effects on Physical, Chemical Properties and Survivability of Probiotic Bacteria During Cold Storage, World Applied Sciences Journal 34(9) :(2016) 1190–1196.
- [43] V. Paul, R. Ezekiel, and R. Pandey. Sprout suppression on potato: need to look beyond CIPC for more effective and safer alternatives, Journal of food science and technology 53(1): (2016) 1-18.
- [44] R.H. Perez, T. Zendo, and K.Sonomoto, Novel bacteriocins from lactic acid bacteria (LAB): various structures and applications, Microbial Cell Factories 13: (2014) Suppl 1:1-13.
- [45] J. O. Pereira, J. Soares, S. Sousa, A. R. Madureira, A. Gomes, and M. Pintado, Edible films as carrier for lactic acid bacteria, LWT 73: (2016) 543-550.
- [46] S. A. Petropoulos, Â. Fernandes, N. Polyzos, V. Antoniadis, L. Barros, and I. CFR Ferreira, The Impact of fertilization regime on the crop performance and chemical composition of potato (*Solanum tuberosum* L.) cultivated in central Greece, Agronomy 10(4) : (2020) 474.
- [47] A.O. Phillips, and G.O. Phillips, Biofunctional behaviour and health benefits of a specific gum Arabic, Food Hydrocolloids 25(2): (2011) 165–169.
- [48] K. Razzaq, A.S. Khan, A.U. Malik, M. Shahidand, and S. Ullah, Role of putrescine in regulating fruit softening and antioxidative enzyme systems in ‘Samar Bahisht Chaunsa’ mango, Postharvest Biology and Technology, 96: (2014) 23-32.
- [49] D. Rico, A.B. Martin-Diana, J.M. Barat, and C. Barry-Ryan, Extending and measuring the quality of fresh-cut food and vegetables: a review, Trends in Food Science and Technology 18: (2007)373-386.

- [50] A. Saha, R. K. Gupta, and Y. K. Tyagi, Effects of edible coatings on the shelf life and quality of potato (*Solanum tuberosum* L.) tubers during storage, *Journal of Chemical and Pharmaceutical Research* 6(12): (2014) 802-809
- [51] L. Trejo-González, A. I. Rodríguez-Hernández, M.del Rocío López-Cuellar, V. M. Juárez-Martínez, and N. Chavarría-Hernández, Antimicrobial pectin-gellan films: effects on three foodborne pathogens in a meat medium, and selected physical-mechanical properties, *CyTA-Journal of Food* 16(1): (2018) 469-476
- [52] M. Vanoli, M.Grassi, M.Buccheri, and A.Rizzolo, Influence of edible coatings on postharvest physiology and quality of Honeydew melon fruit (*Cucumis melo* L. *inodorus*), *Advances in Horticultural Science* 29(2/3): (2015) 65-74
- [53] X. Zheng, L.Ye, T. Jiang, G.Jing, and J. Li, Limiting the deterioration of mango fruit during storage at room temperature by oxalate treatment, *Food Chemistry* 130(2): (2012) 279-285

الملخص العربي

تطبيق الصمغ العربي كأغشية صالحة للأكل في حفظ ما بعد الحصاد وجودة درنات البطاطس

رواء بلال غنام¹ ، شيماء محمد عبد السلام²، عفاف علي أمين¹، مها أمين هويدي²

¹ قسم صحة الطعام- وحدة الميكروبيولوجي- المعهد القومي للتغذية – القاهرة - مصر

² قسم النبات - كلية النبات للآداب والعلوم والتربية -جامعة عين شمس- القاهرة- مصر

الملخص العربي:

تستخدم المواد الغروية على نطاق واسع في صناعة المواد الغذائية، وقد تم استخدامها مؤخرًا وخاصة الصمغ العربي (GA) على نطاق واسع كأغشية / أغلفة قابلة للأكل لتمديد فترة صلاحية الفاكهة والخضراوات بعد الحصاد.

الهدف من هذه الدراسة هو تقييم تأثير غشاء الصمغ العربي (GA) القابل للأكل ممزوجا بالجلسرين و كلوريد الكالسيوم (CaCl₂) كمكونان أساسيان مع جميع أفلام الصمغ العربي وثلاث معاملات مختلفة من GA مع EDTA و *L. paracaseisupernatant* وذلك ع النحو التالي: (GA / EDTA / *L. paracaseisupernatant*) ، (GA / *L. paracaseisupernatant*) و (supernatant / EDTA / *L. paracaseisupernatant*) في الحفاظ على درنات البطاطس المخزنة عند 8 درجات مئوية و 5 ± 30 درجة مئوية لمدة 35 يومًا مقارنة بالشمع (الغلاف البترولي المستخدم كمادة حفظ). التحليلات الفيزيائية والكيميائية بما في ذلك الأس الهيدروجيني ونسبة فقدان الوزن ونسبة المواد الصلبة الذائبة الكلية (TSS) ، وكذلك الفحوصات الميكروبية (العد الكلي لكل من البكتيريا ، الفطريات و الخمائر ومجموعة البكتيريا المعوية).

أظهرت النتائج أن الفيلم المكون من *GA / EDTA / L. paracaseisupernatant* أعطى أفضل النتائج والتي كانت قريبة جدًا لنتائج الغلاف البترولي (الشمع) مقارنة بالدرنات غير المغلفة (الضابطة) والتي غلفت بالأفلام ذات المعاملات الأخرى ، وقد كان ذلك واضحا من خلال انخفاض الحمل الميكروبي ، و تقليل فقدان الوزن ، كذلك تقليل إجمالي المواد الصلبة الذائبة ، مما أدى الى تحكم أفضل في الأس الهيدروجيني وزيادة فترة الصلاحية.